#: <u>I- 10</u> Page: <u>1 of 3</u> Date: 4/201

Date: 4/2012 Revised: 10/05/2018

Abbreviations with Definitions

The following are definitions or abbreviations and terms used in the Orange County Base Hospital Treatment Guidelines and Standing Orders. This list includes abbreviations commonly used in Orange County EMS standing orders, policies, and procedures and is not an inclusive list of all approved medical abbreviations

< Less than.

≤ Less than or equal.

> Greater than.

≥ Greater than or equal.

Indicates a medication or electrical cardiac conversion standing order.

→ Indicates a key step or decision element of a standing order.

AAA: Abdominal Aortic Aneurysm.

Agitated Delirium: Same as Excited Delirium - a medical and behavioral emergency in which an individual is agitated,

irrational, and disregards their own safety (may be related to acute medical or psychological causes)

ALS: Advanced life support (term is often interchanged for paramedic level treatment/actions).

ALS Escort: To transport with paramedic personnel in attendance and ALS equipment on board transport unit.

ALTE: Apparent Life Threatening Event (only in child 3 years-old or less)

AMS: Altered mental status.

ARN: Authorized Registered Nurse (authorized by Orange County EMS)

BH: Base Hospital.

BHC: Base Hospital Coordinator

BHP: Base Hospital Physician on duty at an Orange County Base Hospital.

BLS: Basic life support (term is often interchanged for BLS level treatment/actions).

Bradycardia: Heart rate less than or equal to 60 bpm.

Bradycardia, symptomatic: Heart rate less than 60 and any of the following: systolic blood pressure less than 90

(pediatric less than 80), altered mental status, rales on lung auscultation and signs of CHF,

poor skin perfusion signs.

BRUE: Same as ALTE (Acute Life Threatening Event)

CHF: Congestive heart failure

COPD: Chronic Obstructive Pulmonary Disease.

Approved:

Review Dates: 5/2016; 8/2018 Initial Release Date: 10/05/2018 Final Implementation Date: 10/05/2018 OCEMS copyright © 2018

#: I- 10 Page: 2 of 3 Date: 4/2012

Revised: 4/2012 Revised: 10/05/2018

CPAP: Continuous Positive Airway Pressure.

CVRC: An Orange County Emergency Medical Services designated Cardiovascular Receiving Center.

Diversion: Formal notification of the EMS system through ReddiNet by a hospital that it is not physically or medically

safe for that facility to accept further patients.

ECG: Electrocardiogram.

Abbreviations with Definitions

ET: Endotracheal.

ERC: Emergency Receiving Center, a hospital designated by the Orange County Emergency Medical Services

Agency to receive 9-1-1 basic life support and advanced life support transports.

Excited Delirium: Same as Agitated Delirium - a medical and behavioral emergency in which an individual is agitated.

irrational, and disregards their own safety (may be related to acute medical or psychological causes)

Extremis: Indicates that a patient has abnormal vital signs and is near cardiopulmonary arrest.

High Flow Oxygen: Oxygen by mask at 10 L/min flow; oxygen by nasal canula at 6L/min flow.

Hypotension: Blood pressure < 90 systolic in adult or adolescent, < 80 systolic in pediatric (under age 15 years-old).

IM: Intramuscular.

IO: Intraosseous.

IN: Intranasal.

IV: Intravenous.

IVP: "Intravenous push" or to administer by intravenous injection.

Kg: Kilogram.

MCI: Mass Casualty Incident.

mg: Milligram.

MI: Myocardial infarction or "heart attack".

MICN: Mobile Intensive Care Nurse on duty at an Orange County Base Hospital.

mL: Milliliter.

NS: Normal saline

OD: Overdose

Approved:

Review Dates: 5/2016; 8/2018 Initial Release Date: 10/05/2018 Final Implementation Date: 10/05/2018 OCEMS copyright © 2018

Page: Date:

3 of 3 4/2012

I- 10

Revised: 10/05/2018

Abbreviations with Definitions

ODT:

Oral dissolving tablet (in reference to ondansetron)

PCR:

Patient care record

Pediatric (child): Male/Female less than age 15 years-old

Perfusion:

Circulation of oxygenated blood to tissues

Perfusion, Adequate:

Is determined by good skin signs, appropriate mental status, and palpable radial (brachial in

pediatric) pulses and/or blood pressure appropriate for age.

PEA:

Pulseless electrical activity.

PO:

Orally (per os).

PVAD:

Preexisting vascular access device.

Skin signs:

Refers to blood perfusion of tissues; good skin signs are uniform skin color appropriate to the individual

skin tone, pink nail beds, warm skin to touch; and in pediatrics, good capillary refill (refill in less than 3

seconds)

SL:

Sublingual (under the tongue)

SMR:

Spinal motion restriction

SNRC:

An Orange County Emergency Medical Services designated Stroke-Neurology Receiving Center.

SO:

Standing order, referring to OCEMS standing orders

SQ:

Subcutaneous.

SVT:

Supraventricular tachycardia.

Symptomatic Bradycardia:

Heart rate less than 60 and any of: systolic blood pressure less than 90, altered mental status,

bradycardia: rales on lung auscultation and signs of CHF, poor skin perfusion signs.

TC:

Trauma Center

TG:

Treatment Guideline.

VT:

Ventricular tachycardia.

VF:

Ventricular fibrillation.

Approved:

Review Dates: 5/2016; 8/2018 notto Initial Release Date: 10/05/2018 Final Implementation Date: 10/05/2018 OCEMS copyright © 2018