



**Preparation and Dosing of Push Dose Epinephrine -  
Adult/Adolescent**

**INDICATIONS:**

1. Suspected Septic Shock unresponsive to fluid resuscitation.
2. Distributive (spinal) Shock unresponsive to fluid resuscitation.
3. Cardiogenic Shock unresponsive to initial fluid challenge (250 mL normal saline) or presenting with evidence of pulmonary edema (pulmonary basilar rales).

**CONTRAINDICATIONS:**

1. Hypovolemic Shock (hemorrhage or dehydration) prior to fluid resuscitation and volume replacement.
2. Non-shock (perfusing) states
3. Suspected stimulant drug intoxication

**PROCEDURE:**


**Base Hospital contact and order required.**

**Mixing instructions:**

- Take epinephrine 1 mg of 0.1 mg/mL preparation (cardiac epinephrine) and waste 9 mL of epinephrine.
- Into that syringe, withdraw 9 mL of normal saline from the patient's IV bag. Shake well.
- Mixture now provides 10 mL of epinephrine at a 10 mcg/mL concentration.

**Push Dose:**

- 0.5 mL (5 mcg) IV/IO, every 3 minutes, titrate to a SBP > 90.

Approved: 

Review Date: 02/19  
Final Date for Implementation: 04/01/2019  
OCEMS copyright © 2016