B-040b 1 of 1 Page: Orig. Date: 10/01/2018 Revised:

09/11/2019

INDICATION:

Pulseless cardiopulmonary arrest Adolescent/Adult to provide continuous chest compression.

CONTRAINDICATIONS:

- Chest circumference below or beyond the fit range of the device chest circumference band.
- Recent chest surgery or a chest tube in place.
- Left ventricular assist device in place.

PROCEDURE:

- Remove clothing from thorax area.
- Apply defibrillation-monitor pads to chest in usual location and manner. If device is pass through defibrillator capable, follow manufacturer instructions for specific device.
- Place device next to patient and activate, assuring device powers up and is ready to place on patient.
- Supine position technique:
 - O Supporting head and neck, slide back plate under patient's back.
 - Roll patient from side to side to properly position back plate.
 - O While one rescuer resumes manual chest compressions, have second rescuer open and prepare plunger upper part and attach to back plate nearest rescuer preparing plunger.
 - O Stopping manual chest compression, attach second part of plunger part to back plate.
 - O Assure device is assembled correctly and plunger clicked in locks to back plate.
 - The plunger compression point should be same as for manual chest compression.
 - Using a finger to assess, assure lower edge of plunger is immediately above the end of the sternum.
 - O Adjust plunger so that it contacts the patient's chest without compressing the chest.
 - Activate compressor and assess for positioning and proper functioning.
 - Apply stabilizing strap if supplied with device.
 - When lifting patient to place on gurney, temporarily pause device and reactivate when positioned on gurney (assure plunger is still in proper position).

DOCUMENTATION:

- Document time device placed.
- Check "AutoPulse®" box on PCR.

Carl Schults, MO.

Review Date: 09/2019

Final Date for Implementation: 10/01/2019 OCEMS copyright © 2019