



Notes/Definitions

- Early signs of shock include: tachycardia, delayed cap refill > 2 seconds, pallor, anxious or irritable, tachypnea, cool skin.
- Late signs of shock include: hypotension, unresponsive or somnolent, mottled extremities, worsening tachycardia, tachypnea.

Base Hospital Contact if meets Trauma Criteria

BLS Interventions

- Refer to BLS Standing Orders (SO-B-001).
- Apply cervical collar and /or spinal motion restriction as applicable.

Initial Assessment

xABCDE Assessment: Prioritize life-threatening hemorrhage and shock recognition.

- **(X) Exsanguinating Hemorrhage Control**
 - Immediately apply direct pressure to control bleeding. Scalp hemorrhage can be life threatening.
 - Use of hemostatic dressings indicated for deep wounds or junctional areas.
 - Apply tourniquet 2 to 4 inches above injury in the setting of life-threatening injury.
 - If bleeding is uncontrolled with single tourniquet, apply 2nd tourniquet proximally to the first.
 - Reassess dressings and tourniquets after patient movement.
- **(A) Airway**
 - Ensure patent airway and dislodge any airway obstruction.
 - Ensure proper positioning for adequate ventilation and suction as needed.
 - Consider insertion of nasopharyngeal airway (NPA) or oropharyngeal airway (OPA) per SO-FR-004 and SO-FR-005.
 - Consider BVM for moderate to severe distress, apnea, agonal respirations, or altered mental status with hypoventilation or hypoxia despite maximal supplemental O₂.
- **(B) Breathing**
 - Assess vital signs, including pulse oximetry.
 - Administer supplemental oxygen as necessary to maintain SpO₂ ≥ 95%.
 - Apply vented chest seal to sucking chest wound or large open chest wall injury.
- **(C) Circulation**
 - Aggressively control external hemorrhage.
 - Assess for signs and symptoms of shock:
 - Early signs: tachycardia, delayed cap refill > 2 seconds, pallor, AMS (anxious / irritable), tachypnea, cool skin.



- Late signs: hypotension, AMS (somnolence / unresponsive), mottled extremities, worsening tachycardia and tachypnea.
- **(D) Disability**
 - Apply cervical collar for spinal cord protection.
- **(E) Environmental**
 - Reduce heat loss by removing wet clothing and drying the patient.
 - Cover the patient with warm, dry insulating material covered with a barrier.

ALS Interventions

- Assess Pediatric Assessment Triangle (PAT)
 - General appearance: tone, interactiveness, consolability, look / gaze, speech / cry
 - Work of breathing: retractions, nasal flaring, abnormal airway sounds, positioning
 - Circulation to skin: pallor, mottling, cyanosis
- When BLS airway management cannot maintain adequate ventilation or oxygenation and criteria met, consider:
 - Supraglottic airway insertion for ≥ 50 kg (PR-135).
 - Endotracheal intubation for > 36 kg (PR-030).
- For chest trauma with unilateral decreased or absent breath sounds and **hypoxia**, assess for signs of a tension pneumothorax. Signs include acute respiratory distress, respiratory arrest, circulatory collapse (hypotension, poor perfusion), distended neck veins.
 - If chest seal previously applied remove the dressing, allow air to escape and reapply dressing.
 - **Contact Base Hospital** for an order for needle decompression (needle thoracostomy) (refer to PR-065).
 - **Base Hospital Order** required in all pediatric needle decompressions except in setting of an MCI, remote rescue, tactically unstable scene, or late signs of shock. Provide documentation for these settings if base order not obtained.
- Apply capnography (EtCO₂) as tolerated.
 - Maintain EtCO₂ between 35 and 45 mmHg.
 - If EtCO₂ > 50 , ensure proper positioning for adequate ventilation and suction as needed.
 - Reevaluate need for advanced airway.
- Obtain the patient's estimated weight utilizing an approved pediatric length-based resuscitation tape.
- Obtain IV / IO if applicable.
- If hypotensive (refer to Vital Sign table below), administer:



► **Normal Saline 20 mL / kg IV / IO** (maximum dose 250 mL).

- For continued hypotension, may repeat the same dose twice for total of three boluses.

- Initiate cardiac monitor as needed. Interpret, monitor, and document rhythm.
- Assess GCS. Use pediatric GCS for patients aged ≤ 2 years (refer to B-02).

- Treat for pain choosing one of the following to administer:

► **Fentanyl 2 mcg / kg IN / IV / IM** (maximum single dose 50 mcg IV / IM or 100 mcg IN)

- **Base Hospital Order** required for age < 2 years old.
- Do not exceed a total dose of 100 mcg.
- May repeat once after 3 minutes.
- Hold if hypotensive.

► **Morphine sulfate 0.1 mg / kg IV / IM** (maximum single dose 5 mg)

- **Base Hospital Order** required for age < 2 years old.
- Do not exceed a total dose of 10 mg.
- May repeat once after 3 minutes.
- Hold if hypotensive.

► **Ketorolac 0.25 mg / kg IV / IO / IM** (maximum single dose of 15 mg) once

- *Contraindicated in ages less than 2 years old and designated trauma center patients.*

- For nausea or vomiting administer choosing one of the following to administer:

► **Ondansetron (Zofran) 4 mg ODT** for age > 4 years old with intact airway reflexes once.

► **Ondansetron (Zofran) 0.1 mg / kg IV / IO** for age > 3 months old once.

Transport Considerations

- **Base Hospital Contact required** (CCERC preferred) for patients meeting trauma criteria (refer to OCEMS Policy 310.30 or 310.31). ALS transport to the most appropriate Trauma Center as directed by the Base Hospital.
- ALS providers may determine ALS or BLS level of transport to the nearest, most appropriate ERC for patients not meeting trauma center criteria.
- Pediatric patients are to be transported securely in a size appropriate restraint system (e.g. Neo-mate, Pedi-mate, car seat, etc.) if available.

Additional Considerations

Specific Injury Management

Amputation

- Control hemorrhage as stated above.
- Dress stump with sterile dressing.
- Wrap amputated part in sterile dry material, place in sealed container, then place container in ice.
- Do not allow tissue to directly contact ice or water.



ORANGE COUNTY EMERGENCY MEDICAL SERVICES

Prehospital Standing Orders / Treatment Guidelines

GENERAL INJURY AND TRAUMA - PEDIATRIC

#: P-015

Page: 4 of 5

Orig. Date: 11/1/2016

Revised: 10/01/2026

Evisceration

- Cover with large saline-moistened sterile dressing.
- Do not replace abdominal contents.
- For larger wounds, consider a plastic wrap or occlusive dressing over the moist gauze to maintain moisture and protect from contamination. Avoid packing directly into the wound; instead, control external bleeding while protecting the bowel.

Impaled Objects

- Stabilize in place.
- Remove only if interfering with airway management or CPR.

Orthopedic Injury

- If angulated and pulseless: gently attempt alignment unless resistance or severe pain occurs.
- If angulated with pulse present: splint in position.
- Open fractures: cover with moist sterile dressing; do not reduce.
 - Exception: traction splint for femur fracture can be reduced.
- Document distal neurovascular status before and after splinting. Always note presence or absence of peripheral pulses and sensation.
- May place cold packs over splinted fractures sites for comfort.

Airbag Deployment

- For eye irritation, brush off any powder around upper face.
- Remove contact lenses if present.
- Irrigate with water or normal saline continuously throughout transport.

Eye Injury

- Cover both eyes loosely.
- Avoid pressure on globe.
- Position upright if tolerated and if spinal motion restriction not required.
- Consider Traumatic Brain Injury (TBI) if altered.

Head Injury

- Maintain airway and assist ventilation as needed.
- Provide supplemental oxygen to maintain SPO₂ > 94%.
- Reassess mental status.
- Avoid hypotension or hypoxemia.

Base Hospital Considerations

- Patients meeting trauma triage criteria (OCEMS Policy 310.30 or 310.31) should be routed to the nearest open OCEMS designated Trauma Center.

Approved:

Review Date: 5/16; 7/17; 9/18; 10/19; 11/23; 8/26
Final Date for Implementation: 4/01/2027



ORANGE COUNTY EMERGENCY MEDICAL SERVICES

Prehospital Standing Orders / Treatment Guidelines

GENERAL INJURY AND TRAUMA - PEDIATRIC

#: P-015

Page: 5 of 5

Orig. Date: 11/1/2016

Revised: 10/01/2026

- Trauma victim designation is determined by the Base Hospital.
- For chest trauma with unilateral decreased or absent breath sounds and **hypoxia**, assess for signs of a tension pneumothorax. Signs include acute respiratory distress, respiratory arrest, circulatory collapse (hypotension, poor perfusion), distended neck veins.
 - ▶ **Order a needle decompression** (needle thoracostomy) per OCEMS PR-065:
 - Patient weight $\leq$ 40 kg use 1.25 inch, 16-gauge catheter over needle
 - Patient weight $>$ 40 kg use an adult decompression needle (minimum 2.5-inch length)
- For hemorrhagic shock related to blunt or penetrating trauma not greater than 3 hours, order:
 - ▶ **TXA 15 mg / kg IV / IO** over 10 minutes at a rate of 1 mL per minute (max dose 1 gm) to be given only once.
 - Refer to OCEMS PR-081 for preparation and dosing.
 - Use the 1 mg / 10 mL concentration of TXA only.
 - Notify Trauma Center of the TXA administration.

Cross References:

SO-B-001 BLS Standing Orders

Procedure B-002 Glasgow Coma Scale (Score)

SO-FR-004 Oropharyngeal Airway (OPA)

SO-FR-005 Nasopharyngeal Airway (NPA)

PR-030 Endotracheal Intubation (including EtCO₂)

PR-135 Supraglottic Airway Device Placement – Adult / Adolescent

PR-065 Needle Thoracostomy – Pediatric

PR-081 Tranexamic Acid For Bleeding Management - Pediatric

Policy 310.30 Trauma Triage

Policy 310.31 Trauma Triage Algorithm

Approved:

Review Date: 5/16; 7/17; 9/18; 10/19; 11/23; 8/26

Final Date for Implementation: 4/01/2027